



Disinfectant Wipes

Waterfall Fresh



DESCRIPTION

NCLwipes[™] Disinfectant Wipes clean, disinfect and deodorize in one easy step. NCLwipes[™] Disinfectant Wipes save time and labor and are easy to use. No mixing, No measuring, No diluting. Simply pull a wiper from the canister, wipe the surface and discard. NCLwipes[™] Disinfectant Wipes is a broad spectrum disinfectant that is effective against SARS-Related Coronavirus 2¹ (SARS-CoV-2) in 15 seconds, Norwalk Virus in 10 minutes, and most other viruses, bacteria in 4 minutes.

PRODUCT FEATURES / BENEFITS

- Effective against SARS-Related Coronavirus 2¹ (SARS-CoV-2) in 15 seconds.
- Kills Cold and Flu Virus.*,1
- · Controls the growth of odor-causing bacteria for 72 hours.
- 3 steps in 1- wipes, scrubs stubborn spots and disinfects² so you finish faster.
- Everyday surface¹ disinfectant.
- Kills 99.9% of Bacteria in just 15 seconds.³
- Effective in the presence of 5% serum contamination.
- Eliminates 99.9% of Pseudomonas aeruginosa, Staphylococcus aureus -Methicillin-Resistant, Escherichia coli, Enterococcus faecalis- Vancomycin Resistant, Salmonella enterica in just 4 minutes!
- Leaves surfaces1 clean and disinfected.
- An effective way to clean and deodorize against odor-causing bacteria.
- Faster acting wipe formula.
- A one-step disinfectant² that is effective against a broad spectrum of bacteria when used as directed.
- Phosphate free.
- This product will clean and disinfect hard, non-porous, non-food surfaces.
- Waterfall Fresh Fragrance.
- · Strong wipe for tougher jobs.
- Textured for better scrubbing power.
- · Just pull wipe and toss.
- Recyclable packaging.
- Streak free.
- Reduce risk of cross contamination between treated surfaces.¹
- Ready to Use.
- · Handles daily clean-ups with no need for additional sprays, sponges or buckets.

APPLICATION AREAS

Schools, Colleges, Hospitals, Medical Offices, Nursing homes, Day care centers, Nurseries, Institutional facilities, USDA inspected food processing facilities, Office buildings, Restaurants, Grocery stores, Convenience stores, Hotels, Motels, Prisons, Salons

APPLICATION SURFACES

Chairs, Counters, Desks, Doorknobs, Faucets, Glass, Light switches, Showers, Sinks, Tables, Telephones, Toilets, Diaper changing tables, and other hard nonporous surfaces made of Acrylic, Metal, Plastic, Stainless steel, Upholstery-vinyl and Plastic.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use as a diaper wipe or for personal cleansing. This is not a baby wipe! Do not flush in toilet.

TO CLEAN AND DEODORIZE: Wipe hard, nonporous surface with wipe. Let air dry. Discard used wipe in trash. Do not flush in toilet.

TO CLEAN AND DISINFECT: Wipe hard, nonporous surface with wipe until surface is visibly wet. Use enough wipes to visibly wet surfaces. Use enough wipes to keep surfaces visibly wet for 4 minutes. Allow surface to remain visibly wet for 4 minutes. For Norwalk Virus, Feline Calicivirus, Norovirus and Rotavirus, treated surfaces must remain visibly wet for 10 minutes. If surfaces are visibly dirty, clean first with another wipe before disinfecting. Discard used wipe in trash. Do not flush in toilet.

FOR MOLD AND MILDEW STAIN CONTROL

NCL wipes[™] Disinfectant Wipes will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, nonporous surfaces. Wipe surface and let it remain visibly wet for 4 minutes. Repeat treatment every seven days, or more often if new growth appears.

SANITIZING

To sanitize hard, non-porous, non-food contact surfaces; wipe surface, using enough wipes to allow treated surface to remain visibly wet for 15 seconds for Staphylococcus aureus, Klebsiella pneumoniae, and Campylobacter jejuni. Let surface dry. For visibly soiled surfaces clean visible soil first.

*KILLS HIV, AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in

health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 or HBV or HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Clean-up must always be done wearing protective gloves, gowns, masks and eye protection.

Cleaning procedure: Blood and other body fluids containing HIV or HBV or HCV must be visibly cleaned from surfaces and objects before application of NCL wipes[™] Disinfectant Wipes. NCL wipes[™] Disinfectant Wipes can be used for this purpose.

Contact time: Leave surface visibly wet for 1 minute for HIV-1. Use appropriate contact time as listed on label for all other viruses and bacteria claimed. **Disposal of Infectious Material:** Blood, body fluids, cleaning materials and clothing must be disposed of according to local regulations for infectious waste disposal.

This product is for use on hard nonporous surfaces not conducive to treatment by immersion or excess liquid against Avian Influenza A.

This product is not for use on medical device surfaces.

*.1 On hard, non-porous, non-food surfaces. ² When used according to the disinfectant directions ³ When used according to the sanitization directions



continued on back side





Disinfectant Wipes

Waterfall Fresh



EFFICACY	
Bactericidal Test Results	Contact Time (in min.)
Acinetobacter baumannii	4
Brevibacterium ammoniagenes	4
Campylobacter jejuni	4
Enterobacter aerogenes	4
Enterococcus faecalis	4
Enterococcus faecalis – Vancomycin resistant	4
Escherichia coli	4
Escherichia coli 0157:H7	4
Escherichia coli ESBL (antibiotic resistant)	4
Klebsiella pneumoniae	4
Legionella pneumophilia	4
Listeria monocytogenes	4
Pseudomonas aeruginosa	4
Pseudomonas cepacia – Strain 25416	4
Salmonella enterica	4
Salmonella schottmuelleri	4
Salmonella typhi	4
Serratia marcescens	4
Shigella dysenteriae	4
Staphylococcus aureus	4
Staphylococcus aureus – Multi-drug resistant	4
Staphylococcus aureus – MRSA Staphylococcus aureus – CA-MRSA (USA 300/400)	-
Staphylococcus aureus – VISA	4
Streptococcus pyogenes	4
Vibrio cholerae	4
Virucidal Test Results	Contact Time (in min.)
*Hepatitis B Virus (HBV)	4
*Hepatitis C Virus (HCV)	4
*Herpes Simplex Virus Type 1	4
*Herpes Simplex Virus Type 2	4
*HIV-1 (AIDS Virus)	1
*Human Coronavirus	4
*Influenza Type A Virus / Brazil (the Flu Virus)	4
*Influenza A Virus H1N1	4
*2013 Influenza A virus (H7N9)	4
*Norwalk Virus (Norovirus)	10
*Respiratory Syncytial Virus	4
*Rotavirus	10
*SARS associated Coronavirus	4
*SARS related Coronavirus 2	15 (sec.)
*Vaccinia Virus	4
Virucidal Test Results (Animal)	Contact Time (in min.)
Avian Influenza Virus (H3N2)	4
Avian Influenza Virus (H5N1)	4
Canine Distemper Virus	4
Feline Calicivirus	10
Newcastle's Disease Virus	4
Pseudorabies Virus Fungicidal Test Results	Contact Time (in min.)
Candida albicans	

Respiratory illnesses attributable to Pandemic 2009 H1N1 are caused by influenza A virus. This product is a broad-spectrum hard surface disinfectant that has been shown to be effective against influenza A virus tested and listed on the label and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).

SPECIFICATIONS

Wipe Size	
Wipes per Canister	
Canisters per Case	
Net Wt / Canister	
Color	
Fragrance	Lemon Fresh
рН	
Flammability	Not Flammable
Shelf Life	1 year minimum in original unopened container

SAFETY INFORMATION

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

Caution. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

8

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor or going for treatment.

STORAGE AND DISPOSAL

Store in original container in areas inaccessible to children. Keep securely closed. Nonrefillable container. Do not reuse or refill this container. Discard empty container in trash or recycle.

*,1 On hard, non-porous, non-food surfaces.

² When used according to the disinfectant directions

³ When used according to the sanitization directions



Candida albicans

401 N. 10th Street · Philadelphia, PA 19123 USA Toll Free: (800) 628-2436 Phone: (215) 922-1200 · Fax: 215-922-5517 Website: www.nclonline.com Contact E-mail: info@nclonline.com