

How to use CITRUS FLOWER-QUAT™



READY-TO-USE LABEL (Diluted 1:32)
EPA REG. NO. 10324-157-0296-E • EPA EST. NO. 2296-94-1

ACTIVE INGREDIENTS:
Dimethyl ammonium chloride 1.27%
Alkal (50% Ca, 40% Cu, 10% Ca) 0.85%
Dimethyl benzoyl ammonium chloride 0.85%
OTHER INGREDIENTS: 22.962%
TOTAL: 100.00%

The product in this container is diluted as directed on the pesticide product label.

DIRECTIONS FOR USE: Follow directions for use on the pesticide label when applying this product. Pre-clean heavily soiled areas. Apply use solution of 4 oz. of this product per gal. of water to disinfect hard, non-porous surfaces with a sponge, brush, cloth, mop, by immersion, auto scrubber, hand pump trigger spray device, coarse trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Treated surfaces must remain wet for 10 minutes. Wipe dry with a clean cloth or allow to air dry. Rinsing of floors is not necessary unless they are to be waxed or polished. Prepare a fresh solution daily or when visibly dirty.

PRECAUTIONARY STATEMENTS
CAUTION: KEEP OUT OF REACH OF CHILDREN.
Causes mild skin irritation. Causes eye irritation.
PREVENTION: Wash thoroughly after handling. Wear protective gloves. Wear eye/face protection.
IF IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
Get medical attention if irritation develops and persists.
IF ON SKIN OR CLOTHING: Remove contaminated clothing. Wash with plenty of soap and water. Get medical attention if irritation develops and persists. Wash contaminated clothing before reuse.
IF SWALLOWED: Rinse mouth. Get medical attention if irritation develops and persists.
IF INHALED: Move to fresh air. Get medical attention if irritation develops and persists.

World Class Cleaning & Hygiene Solutions
401 N. 10th Street • Philadelphia, PA 19123
(800) 922-1200 • (215) 922-1200 • www.nclonline.com

DUAL-BLEND EQUIPMENT DIRECTIONS:

Chemical Selector Dial

1. Turn the chemical selector dial to the desired chemical position, indicated by the directional arrow.
2. Dispense chemical by either activating the bottle fill lever or bucket fill button.
3. For rinse water, point the dial straight down at the water symbol. This is not potable water, do not drink.

Bottle Fill

1. Slide the bottle tube inside a the spray bottle and raise the bottle to engage the u-shaped lever of the bottle fill assembly.
2. To stop filling, lower the bottle and the lever will disengage.
3. Be careful when removing the spray bottle so as to not splash residual chemical.

Bucket Fill

1. To start the flow of diluted chemical, press button on unit above selector dial.
2. For continuous flow, twist button clockwise.
3. To disengage, release button or twist button counterclockwise.


NOTE: The DUAL-BLEND® bottle is designed for use with the DUAL-BLEND® Wall Dispenser and the DUAL-BLEND® Portable Dispensing Unit. Dilution tips built into the product containers provide precise, accurate dilution for every product, every time, regardless of dispensing equipment chosen.

DIRECTIONS FOR USE:

FOR USE AS A GENERAL, HOSPITAL, MEDICAL DISINFECTANT, FUNGICIDE, VIRUCIDE*, CLEANER: Pre-clean visibly soiled areas. Apply use solution of 4 oz. of this product per gal. of water to disinfect hard, non-porous, non-food contact surfaces with a sponge, brush, cloth, mop, by immersion, auto scrubber, hand pump trigger spray device, coarse trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Treated surfaces must remain visibly wet for 2 minutes to kill SARS-CoV-2 or for 10 minutes to kill all organisms listed on the label. Wipe dry with a clean cloth or allow to air dry. Rinsing of floors is not necessary unless they are to be waxed or polished. Prepare a fresh solution daily or when visibly dirty.

***KILLS HIV, HBV AND HCV ON PRE-CLEANED HARD, NON-POROUS SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of hard, non-porous surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1), Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS. Personal Protection: Wear protective latex gloves, gowns, masks and eye protection. Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from hard, non-porous surfaces and objects before application of this product. Disposal of Infectious Materials: Blood and other body fluids, cleaning materials and clothing must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal. Contact Time: Allow hard, non-porous surfaces to remain visibly wet



for 2 minutes to kill HIV and 10 minutes for all organisms listed on the label.

NON-FOOD CONTACT SURFACE SANITIZING: Pre-clean visibly soiled areas. Add 4 oz. of this product per gal. of water. Apply solution to hard, non-porous surfaces with a sponge, brush, cloth, mop, by immersion, hand pump trigger spray device, coarse trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Treated surfaces must remain visibly wet for 3 minutes. Prepare a fresh solution daily or when visibly dirty.

TO CONTROL MOLD/MILDEW: Pre-clean visibly soiled areas. Prepare use solution by adding 4 oz. of this product per gal. of water. Apply use solution to hard, non-porous surfaces which will effectively inhibit the growth of mold and mildew and their odors. Repeat treatment every seven days, or more often if new growth appears.


NON-ACID TOILET BOWL AND URINAL DISINFECTANT/CLEANING DIRECTIONS: Remove visible soil prior to disinfection.

TO CLEAN WATERFREE/WATERLESS URINALS: Remove any debris from the urinal. Spray 4 - 8 oz. of use solution onto urinal surface. To prepare use solution: Add 4 oz. of this product per gal. of water. DO NOT spray product directly onto cartridge. Wipe surface to clean. Change cartridge as needed. The unit is ready for use.

GENERAL DEODORIZATION: To deodorize, apply 4 - 8 oz. of this product per gal. of water to hard, non-porous surfaces. Wipe up excess liquid with a paper towel or allow to air dry. Rinse with potable water after use on surfaces that come in contact with food.

DISPOSE OF UNUSED PRODUCT IN ACCORDANCE WITH LOCAL, STATE AND FEDERAL REGULATIONS - PLEASE RECYCLE CONTAINER

PPE INFORMATION: Respiratory protection is generally not required. No special ventilation is normally required. The use of safety glasses and gloves is recommended. No other protective equipment is required.



Philadelphia, PA 19123 • 1-800-NAT-CHEM • (215) 922-1200
E-mail: info@nclonline.com • Web Site: www.nclonline.com